Manuscript for *Revista Mexicana de Astronomía y Astrofísica* (2007)

arXiv:1209.0290v1 [astro-ph.SR] 3 Sep 2012

LONG-TERM PHOTOMETRIC AND SPECTROSCOPIC OBSERVATIONS OF THE NEAR-CONTACT BINARY KR CYGNI[2]

E. Sipahi,[1] C. İbanoğlu[1] and Ö. Çakırlı,[1]

Draft version: September 4, 2012

RESUMEN

ABSTRACT

We present the multi-color, five-year light curves and the first radial velocities of the near-contact binary system KR Cyg. We derived the masses of the components as 2.88±0.20 M$_\odot$ and 1.26±0.07 M$_\odot$ and the radii as 2.59±0.06 R$_\odot$ and 1.80±0.04 R$_\odot$. Analyses of the UBVR light curves and the radial velocities indicate that none of the components exactly fill their corresponding Roche lobes. We have calculated the distance to the system of KR Cyg as 411±12 pc using the observed apparent UBV magnitudes and the bolometric corrections for the component stars. We also searched for the empirical determination of albedo and effective temperature of the cooler, less massive star of KR Cyg, and of two similar near contact binaries AK CMi, and DO Cas. The residuals between the observed and computed fluxes are **attributed to the effect of mutual illumination which heats the surface layers of the illuminated star and does vary not only its bolometric albedo but also its limb-darkening coefficient and gravity-brightening exponent.** The analysis of the light curves shows that the effective albedos are generally smaller than that expected from an envelope of convective star, being mostly departed from the theoretical value at the B passband. As the reflected light diminishes the effective temperature and, therefore, the luminosity of the irradiated star increase. The observed bluer U-B colors during primary minimum are attributed to the effects of mutual irradiation and multiple scattering processes which may alter several characteristics of these systems.

Key Words: stars: binaries: close — binaries: eclipsing — binaries: general — binaries: spectroscopic — stars: individual: KR Cyg, AK CMi, DO Cas

1. INTRODUCTION

The light variability of KR Cyg (HD 333645, $V = 9^m.19$) was discovered by Schneller (1931a,b,c). The light curves obtained by photographic observations were published by Nekrasova (1945) and Wachmann (1948) and the visual observations by Lassovzsky (1936), Gaposchkin (1953) and Tsesevich (1954). Both visual and photographic observations showed that the light variations were originated from the eclipses. The light curves were very similar to those of Algol-type binaries. The first photoelectric observations were made by Vetesnik (1965). He obtained blue and yellow light curves of the system. The passbands used in the observations were very similar to those Johnson's B and V filters. A preliminary solution of the light curves leads him to the conclusion that the system consists of a B9 and an F5 main-sequence stars. He classified the system as a β Lyrae type by visual inspection of the light curves. He also called attention some peculiarities in the course of the light curve. Later on Vetesnik's two-color light curves were analysed by Wilson & Rafert (1980) using a contemporary method, i.e. the Wilson-Devinney code (Wilson & Devinney 1971), henceforth WD. The main difference between the two solutions is the primary star's radius which is fifteen per cent larger in the later than that found in the former analysis. In addition, Wilson & Rafert (1980) use an effective temperature for the primary star corresponding to a B7 star. The observations in the primary minimum of Vetesnik's light curves were re-analysed by A-Naimiy et al. (1985) using the Fourier techniques in the frequency domain. They find that, in contrary to the older estimates, more massive star has smaller radius than its less massive companion.

---

[1] Ege University, Science Faculty, Department of Astronomy and Space Sciences, 35100 Bornova, İzmir, TURKEY

[2] BASED ON OBSERVATIONS COLLECTED AT TÜBİTAK NATIONAL OBSERVATORY (ANTALYA, TURKEY).

Sipahi & Gulmen (2004) published three-color photometric observations and the resultant light curves of KR Cyg. The light curves were analysed individually and simultaneously by the WD code. They determined a photometric mass-ratio to be 0.43 and suggested that the cooler, less massive component is nearly filling its corresponding Roche lobe. Recently, one of us (Sipahi 2012) collected all available times of light minimum and studied orbital period of the system. She arrived at a result that the differences between the times of observed and calculated (O-C) show a periodic oscillation with an amplitude of 0.001 days and a period of about 76 years. This periodic change was attributed to a hypothetical third component.

Shaw (1990) defined a group of close binary stars named near-contact binaries (NCB) in which both components fill or nearly fill their critical Roche lobes. KR Cyg is included in the list of NCBs. According to his definition the stars of the near-contact binaries are near enough to each other to have strong proximity effects like W UMa type systems but are not in contact. As it is known the mass and energy exchange are being occurred between the components in the contact systems. This energy exchange is resulted in both components to come to nearly the same effective temperature. The main difference between the near-contact and contact binaries is the non-existence of the energy exchange through the neck of material connecting the components in the near-contact systems. The non-existence of energy exchange in the case of NCB allows a large temperature difference between the components. Some NCBs show photometric asymmetries such that the system is brighter at the first quarter than that at the second one. This

TABLE 1

THE COORDINATES, APPARENT VISUAL MAGNITUDES AND THE SPECTRAL TYPES OF THE STARS OBSERVED.

Star	Alpha	Delta	V (mag)	Sp
KR Cyg	20 09 06	30 33 01	9.23	A0V
HD 191398	20 08 39	30 20 15	9.01	A0V
HD 333664	20 09 18	30 13 39	9.58	A0

asymmetry is attributed to the primary star being one hemisphere, facing at phase about 0.25, is hotter than that at phase 0.75, or some material is obscuring the surface of the primary visible at phase 0.75. Therefore the NCBs are assumed to be lying in key evolutionary states of close binary systems. Shaw (1994) renewed the list of NCBs in which over 130 systems are given. He proposes that the NCBs are precursors of the A-type W UMa contact binaries.

The main aim of this study is to analyse the 1999 and 2000 BVR, and 2003, 2004, and 2005 UBVR light curves and the first radial velocities obtained by us. We derive the absolute physical parameters of the components and discuss light curve peculiarities by taking into account the mutual irradiation and therefore the changing in the value of albedo of the irradiated star for four passbands.

2. OBSERVATIONS

2.1. *Photometric observations*

The observations were made with the High-Speed Three-Channel Photometer attached to the 48 cm Cassegrain telescope at Ege University Observatory. HD 191398 and HD 333664, respectively, are taken as the comparison and check stars. These stars are nearly the same apparent magnitude and spectral type with the variable star. Moreover, they are very closely located with the variable on the plane of sky which make the effect of the atmospheric extinction on the differential magnitudes almost negligible. In Table 1 the coordinates, apparent visual magnitudes and spectral types of the variable and the comparison stars are given. The observations were made using the BVR passbands in 1999 and 2000, and UBVR passbands in 2003, 2004 and 2005. Though the comparison star is so close to the variable, the differential magnitudes, in the sense variable minus comparison, are corrected for atmospheric extinction. The nightly extinction coefficients were derived from the observations of the comparison stars in each passband.

2.2. *Spectroscopic observations*

Optical spectroscopic observations of KR Cyg were obtained with the Turkish Faint Object Spectrograph Camera (TFOSC) attached to the 1.5 m telescope on 4 nights (July-August, 2011) under good seeing conditions. Further details on the telescope and the spectrograph can be found at http://www.tug.tubitak.gov.tr. The wavelength coverage of each spectrum was 4000-9000 Å in 12 orders, with a resolving power of $\lambda/\Delta\lambda$ 7 000 at 6563 Å and an average signal-to-noise ratio (S/N) was ∼120. We also obtained a high S/N spectrum of the α Lyr (A0 V) and HD 27962 for use as templates in derivation of the radial velocities (Nidever et al. 2002).

The electronic bias was removed from each image and we used the 'crreject' option for cosmic ray removal. Thus, the resulting spectra were largely cleaned from the cosmic rays. The echelle spectra were extracted and wavelength calibrated by using Fe-Ar lamp source with help of the IRAF ECHELLE (Tonry & Davis 1979) package.

The stability of the instrument was checked by cross correlating the spectra of the standard star against each other using the FXCOR task in IRAF. The standard deviation of the differences between the velocities measured using FXCOR and the velocities in Nidever et al. (2002) was about 1.1 km s^{-1}.

3. ANALYSES

3.1. *Radial velocities*

To derive the radial velocities for the components of binary system, the four TFOSC spectra of the eclipsing binary, obtained near the quadratures, were cross-correlated against the spectrum of α Lyr on an order-by-order basis using the FXCOR package in IRAF. The majority of the spectra showed two distinct cross-correlation peaks in the quadrature, one for each component of the binary. Thus, both peaks were fitted independently in the quadrature with a *Gaussian* profile to measure the velocity and errors of the individual components. If the two peaks appear blended, a double Gaussian was applied to the combined profile using *de-blend* function in the task. For each of the four observations we, then, determined a weighted-average radial velocity for each star from all orders without significant contamination by telluric absorption features. Here we used as weights the inverse of the variance of the radial velocity measurements in each order, as reported by FXCOR.

We adopted a $two - Gaussian$ fit algorithm to resolve cross-correlation peaks near the first and second quadratures when spectral lines are visible separately. Fig. 1 shows examples of cros-correlations obtained by using the largest FWHM at nearly first and second quadratures. The two peaks, correspond to each component of KR Cyg. The stronger peaks in each CCF correspond to the more luminous component which has a larger weight into the observed spectrum.

The heliocentric RVs for the primary (V$_p$) and the secondary (V$_s$) components are listed in Table 2, along with the dates of observation and the corresponding orbital phases computed with the new ephemeris given by Sipahi (2012). The velocities in this table have been corrected to the heliocentric reference system by adopting a radial velocity of 9.5 km s^{-1} for the template star α Lyr. The RVs listed in Table 2 are the weighted averages of the values



Fig. 1. Sample of CCFs between KR Cyg and the RV template spectrum around the first and second quadrature.

TABLE 2

HELIOCENTRIC RADIAL VELOCITIES OF KR CYGNI. THE COLUMNS GIVE THE HELIOCENTRIC JULIAN DATE, THE ORBITAL PHASE, THE RADIAL VELOCITIES OF THE TWO COMPONENTS WITH THE CORRESPONDING STANDARD DEVIATIONS.

HJD 2400000+	Phase	Star 1		Star 2	
		V_p	σ	V_s	σ
55751.36069	0.8080	73.1	11.8	-290.8	11.3
55796.38592	0.0826	-92.3	7.8	77.7	11.2
55835.36321	0.2013	-150.4	6.8	188.3	9.5
55835.46128	0.3173	-144.6	9.9	184.1	9.9



Fig. 2. Radial velocities of the components.

obtained from the cross-correlation of orders #4#, #5#, #6# and #7# of the target spectra with the corresponding order of the standard star spectrum. The weight $W_i = 1/\sigma_i^2$ has been given to each measurement. The standard errors of the weighted means have been calculated on the basis of the errors (σ_i) in the RV values for each order according to the usual formula (e.g.Topping (1972)). The σ_i values are computed by FXCOR according to the fitted peak height, as described by Tonry & Davis (1979).

The measured radial velocities of both stars are plotted in Fig. 2 against the orbital phase. First we analysed the radial velocities for the initial orbital parameters. We used the orbital period held fixed and computed the eccentricity of the orbit, systemic velocity and semi-amplitudes of the RVs. The results of the analysis are as follows: e=0.001±0.001, i.e. formally consistent with a circular orbit as expected, γ= -43±3 km s^{-1}, K_1=110±4 km s^{-1} and K_2=251±7 km s^{-1}. Using these values we estimate the projected orbital semi-major axis and mass ratio as: $a\sin i$=6.028±0.127 R$_\odot$ and $q = \frac{M_2}{M_1}$=0.438±0.017.

3.2. *Effective temperature of the primary star*

The spectral type of a star can be derived from the multi-color photometry and spectroscopy, or both. Unfortunately, UBV measurements of the system are not available. Therefore we observed the system with the UBV standard stars some nights and obtained the standard apparent visual magnitude and colors at mid-secondary eclipse as: V=9.231±5, U-B=0.101±15, and B-V=0.156±7 mag. The Q-parameter of the Johnson's wide-band photometry, defined as Q=(U-B) -0.72 (B-V) (Johnson & Morgan 1953), is independent of interstellar reddening for early type stars. Using the observed (U-B) and (B-V) colors at secondary eclipse we calculated the Q-parameter as -0.011 for the hotter star. Then, we estimated the intrinsic color of $(B\text{-}V)_0$ = -0.021 mag from the tables given by Hovhannessian (2004). Since the light contribution of the secondary component does not exceed a few hundreths of magnitudes in the B and V passbands its contribution to the B-V color at maximum light may be ignored. This intrinsic color corresponds to an A0±1 main-sequence star. The average effective temperature deduced from the calibrations of Bohm-Vitense (1981), de Jager & Nieuwenhuijzen (1987), Flower (1996) and Drilling & Landolt (2000) is about 9 810 K. The spectral-type uncertainty leads to an uncertainty of about 150 K in the effective temperature of the primary star. The difference between the observed and intrinsic B-V color, i.e. the interstellar reddening, is about 0.177 mag.

KR Cyg was observed by Hilditch& Hill (1975) in the intermediate-band photometric system. One of their observations falls into the primary eclipse. Excluding this observation we calculated the colors of $u-b$=1.555±0.011, $b-y$=0.111±0.010 and c=0.970±0.022 mag. Taking E$(b-y)$=0.73E(B-V) we computed the intrinsic color of $(b-y)_0$=-0.018±0.010 mag which corresponds to an A0 star. The infra-red magnitudes J=9.278±0.022, H=9.169±0.022 and K=9.098±0.018 are given in the 2$MASS$ catalog (Cutri 2003). The infrared colors are calculated as J-H=0.109±0.031 and H-K=0.071±0.028 which correspond to a reddened B9±2 star. The spectral type of the primary star obtained both intermediate and infrared photometric systems is consistent what we derived from the UBV photometry.

3.3. *Analyses of the light curves*

As we stated in §2.1 we have multi-color light curves of KR Cyg obtained with the same instrumentation. We used the recent version of the eclipsing binary light curve modelling algorithm of Wilson & Devinney (1971) (with updates), as implemented in the PHOEBE code of Prša & Zwitter (2005). The code needs some input parameters, such as the effective temperature of the hotter star and the mass-ratio of the system. These two parameters are the key-parameters for the solution which should be estimated before beginning the light curve modelling. The effective temperature of the hotter star, eclipsed in the primary minimum, has been estimated using the photometric indices as 9 810±150 K. The mass ratio, $q=M_2/M_1$, is very important parameter in the light curve analysis, because the WD code is based on Roche geometry which is sensitive to this quantity. The mass ratio of 0.438±0.017 determined from the radial velocities was kept as a fix value. A preliminary estimate for the effective temperature of the cooler component is made using the depths of the eclipses.

The logarithmic limb-darkening coefficients, x_1 and x_2, and bolometric limb-darkening coefficients, x_{bol} and y_{bol}, were determined from tables by Van Hamme (1993) for the primary and secondary components, respectively, taking into account the effective temperatures and the wavelengths of the observations. Standard values of bolometric albedos (Rucinski 1973), and the gravity-darkening coefficients (Lucy 1967) for radiative and convective envelopes were used. The rotational velocities of the components are assumed to be synchronous with the orbital one. The adjustable parameters in the light curves fitting were the orbital inclination i, the surface potentials Ω_1 and Ω_2, the effective temperature of secondary T_2, the luminosity of the primary star L_1.

Using a trial-and-error method, we obtained a preliminary set of parameters which represent the observed UBVR light curves. The code was set in Mode–2, a detached configuration, with coupling between luminosity and effective temperature and no constraints on the potentials. We used a simple reflection treatment as $MREF = 1$ and $NREF = 1$ at the beginning of iterations.

The iterations were carried out automatically until convergence and a solution was defined as the set of parameters for which the differential corrections were smaller than the probable errors. After a few iterations convergence was obtained. The final results of the five-year multi-color light curves' simultaneous analysis of KR Cyg are listed in Table 3. r_1 and r_2 in this table are the mean fractional radii of the hotter and cooler components, respectively. The uncertainties assigned to the adjusted parameters are the internal errors provided directly by the Wilson-Devinney code. We also ran models for KR Cyg in mode Mode–5, the secondary component filling its Roche lobe, in Mode–4, the primary filling its Roche lobe and in Mode–3, both components fill their Roche lobes. Convergence is obtained in Modes–4 and 3 but with larger sum of squared residuals.

The computed light curves (continuous line) are compared with the observations in four passbands in Figure 3. While there is a satisfactory fit between the observed and computed lights for the U passband the binary model does not represent the observed light curves successfully in the B, V, and R passbands. The residuals, in the sense observed minus calculated, are plotted in Fig.4 for a better visibility. The differences between the model and the observed light curves are clearly seen at both maxima in three passbands and particularly in the secondary eclipse in B and R passbands. As pointed out by Shaw (1990) all of the NCBs do not show O'Connell effect, one maximum is higher than the other. If a NCB shows an asymmetry in the light curve, the first maximum is brighter than the second maximum. This difference is generally attributed to the secondary component, in which the visible surface at phase 0.25 is brighter than the other half. While the asymmetries reach to 5% of the total light, the secondary components in these systems contribute less than 5% of the system's total light. Therefore, it is suggested that the visible hemispheres of the hotter components at the orbital phase of 0.75 are either cooler or obscured by a cool material.

In the case of KR Cyg the light curve is almost symmetric in all passbands. At phases of about 0.15 and 0.85 the observed fluxes are larger than those computed in the BVR passbands. In contrary, the observed fluxes are smaller



Fig. 3. The observed five-year UBVR flux scale light curves and computed light curves for KR Cyg.

TABLE 3

FITTED PARAMETERS FOR THE FIVE-YEAR MULTI-COLOR LIGHT CURVES OF KR CYG.

Parameter	U	B	V	R
λ (Å)	3500	4300	5500	6700
N_{obs}	1752	2825	2804	2825
x_1	0.620	0.714	0.611	0.773
x_2	0.871	0.844	0.773	0.682
g_1	1.0	1.0	1.0	1.0
g_2	0.32	0.32	0.32	0.32
A_1	1.0	1.0	1.0	1.0
A_2	0.5	0.5	0.5	0.5
T_1 (K)	9810	9810	9810	9810
$q=M_2/M_1$	0.438	0.438	0.438	0.438
$L_1/(L_{1+2})$	0.9791±0.0005	0.9727±0.0004	0.9467±0.0005	0.9179±0.0006
$i(^\circ)$	87.00±0.02			
T_2 (K)	5 580±10			
Ω_1	2.8645±0.0012			
Ω_2	2.8111±0.0008			
r_1	0.4297±0.0002			
r_2	0.2989±0.0002			
χ^2	1.821			

than those computed at the secondary eclipse in the B, V and R passbands. The components in the NCBs are nearly touch each other but their photosphere are not in physical contact. A gaseous stream and an energy exchange between the stars may be excluded.

3.4. *Search for the empirical derivation of the albedo*

The value of the local emergent bolometric flux and energy balance in a distorted star are significantly dependent upon the assumed values of the gravity darkening exponent and the bolometric albedo (Rafert & Twigg 1980). The reflection effect is used to describe the mutual irradiation of the facing hemispheres of the components in a close binary system. For the stars of intermediate temperatures of about 10 000K the incoming radiation is almost completely absorbed by atoms and ions in the irradiated atmosphere which alters the temperature structure and causes an increased emergent flux. For the stars of convective envelopes half of the incoming radiation is converted to bulk motions of the atmosphere gases (Rucinski 1969). The theoretical value of bolometric albedo is assumed to be about 0.5 for the stars having convective atmospheres. Furthermore, Vaz & Nordlund (1985) calculated bolometric reflection albedo for a particular reflecting main-sequence star with a temperature of 4 500 K and presented the results as a function of angle of incidence and relative incident flux. As pointed out by Cranmer (1993) the effect of mutual irradiation will be strongest on the line-of-centers points of the stars in a close binary system. The stars themselves iterate this effect between one another, e.g., hotter star heats up the cooler that in turn affects the illumination back on the hotter star, and so on. The facing hemispheres of both components are heated up by multiple scattering process.

Therefore we attempted to estimate the values of g_2 and A_2 from analyses of well-defined multi-color light curves of the system. WD code allows us to make parallel solutions by solving any number of subsets of adjustable parameters in the same computer run. By adopting the theoretical values of the less sensitive parameters of limb-darkening coefficients, gravity-darkening exponents and the limb-darkening coefficients we derived the basic set of parameters i, Ω_1, Ω_2, L_1, L_2 and T_2 from the analyses of multi-color light curves. The components of the system KR Cyg are nearly contact and they rotate faster than 100 km s^{-1}, and have a large temperature difference of about 4 200 K. Therefore, the gravity-darkening exponent and the bolometric albedo for the components are expected to be different from the non-rotating and gravitationally undistorted stars. In the second step, we take gravity-darkening exponent, effective temperature, luminosity and effective albedo for the cooler component as adjustable parameters in the analysis of U, B, V, and R light curves separately. A few iterations indicate that there is no significant deviation in the gravity-darkening exponent between the theoretical and calculated values. The iterations are carried on excluding the gravity-darkening. After a few iterations a convergence is attained. The computed values of A_2,and T_2 are given with their standard deviations in Table 5. The



Fig. 4. Residuals, in light units, for the two solutions of KR Cyg's UBVR light curves from top to bottom: A_2=0.5 (fixed, left panels) and A_2 is an adjustable parameter (right panels).

TABLE 4

ABSOLUTE PARAMETERS OF KR CYG.

Parameter	Primary	Secondary
a ($R_\odot$)	6.04±0.13	
V_γ (km s^{-1})	-43±5	
q	0.438±0.017	
Mass ($M_\odot$)	2.876±0.198	1.261±0.074
Radius ($R_\odot$)	2.594±0.055	1.804±0.038
log g (cgs)	4.069±0.012	4.026±0.012
T_{eff} (K)	9 810±150	5 580±150
$(vsin\ i)_{obs}$ (km s^{-1})	141±7	108±9
$(vsin\ i)_{calc.}$ (km s^{-1})	155±3	108±2
log $(L/L_\odot)$	1.750±0.032	0.454±0.050
d (pc)	411±12	
U, B, V (mag)	9.488±0.014, 9.387±0.005, 9.231±0.005	
J, H, K_s (mag)	7.159±0.018, 7.221±0.027, 7.249±0.027	

TABLE 5

THE CALCULATED EFFECTIVE ALBEDOS AND EFFECTIVE TEMPERATURES FOR THE COOLER COMPONENTS OF KR CYG, AK CMI AND DO CAS.

Star	Parameter	U	B	V	R
KR Cyg	A_2	0.435±0.023	0.225±0.016	0.241±0.012	0.304±0.012
KR Cyg	T_2	5600±27	5985±14	5653±12	5336±12
KR Cyg	L_2	0.022	0.041	0.057	0.070
AK CMi	A_2	0.457±0.041	0.132±0.038	0.219±0.028	—
AK CMi	T_2	5658±60	5880±45	5677±32	—
DO Cas	A_2	0.275±0.158	0.377±0.127	0.333±0.089	—
DO Cas	T_2	5441±147	5615±105	5427±74	—

computed L_2 values are also presented. Following the formulation given by Cranmer (1993) and Claret (2001) we connect the bolometric albedo with the intrinsic and incident flux by

$$\sigma T_h^4 = \sigma T_{eff}^4 + F_{incident} \tag{1}$$

$$A_{F_r} = \sigma T_h^4 - T_{eff}^4 \tag{2}$$

where T_{eff} is the effective temperature of the star without external flux, T_h is the effective temperature after irradiation, F_r is the external radial flux and A is the bolometric albedo. The external flux F_r is depended on apparent radius, effective temperature of the irradiating star, and the cosine of the angle of incidence. The amount (1-A)F_r is absorbed and the atmosphere of the irradiated star re-emit this excess energy later. Moreover, Kirbiyik & Smith (1976) and Kirbiyik (1982) proposed that not only the irradiated hemisphere of the star is heated in close binaries. The horizontal fluxes resulting from the incident radiation field can cause large circular currents over the irradiated stellar surfaces and may penetrate into the non-directly irradiated hemisphere. As a result the irradiated star would increase its temperature and luminosity. The values given in Table 5 confirm this theoretical prediction. The empirically derived values of the A_2, and T_2 from the light curves are plotted versus the wavelengths of the observations in Fig. 5. The derived values of A_2 are systematically smaller than their theoretical values of about 0.5. Moreover, the deviations appear to be depended upon the wavelengths. The largest deviation occurs about λ4350. While the albedo gets smaller the effective temperature of the illuminated star increases. The computed value of albedo in the U-passband is close to the theoretically expected value for convective envelopes. The light contribution of the secondary star to the total light is too low for this passband. **The influence of the mutual illumination in close binary systems is not only seen on the bolometric albedos but also in other properties of the stars such as limb-darkening coefficients and gravity-darkening exponents. The reflection effect has a strong influence on limb-darkening coefficients, as already noticed by Vaz & Nordlund (1985), Nordlund & Vaz (1990), Claret & Gimenez (1990). Later on, the effect of illumination on the limb-darkening coefficients is studied numerically by Alencar & Vaz (1999) using the Uppsala Model Atmosphere code in convective line-blanketed atmospheres. Their results show that the limb-darkening coefficients of illuminated atmospheres are significantly different from the non-illuminated ones. They have concluded that the limb-darkening coefficients vary depending on the characteristics of external illumination such as the incidence angle, the amount of infalling flux, the temperature of the illuminating star. In addition, Alencar et al. (1999) showed that external illumination changes the gravity-brightening**

exponent of an illuminated star. They have proposed that the contact and semi-detached eclipsing binary systems should be better represented by illuminated atmospheres, since their components are closer to each other. Their numerical calculations show that the external illumination increases the values of gravity-brightening exponent roughly in proportional to the amount of the incident flux. The classical value of 0.32 for the convective atmospheres may be too small for binary systems with close components.

We also solved the UBV light curves of AK CMi, a similar system to KR Cyg, obtained by Samec et al. (1998). They estimated an effective temperature of 8 510 K for the hotter star. We derived a value of 10 150 K using the UBV measurements of the system at four orbital phases, at maxima and in mid-eclipses. Applying the same procedure as in KR Cyg we obtained A_2, and T_2 for the cooler component. They are listed in Table 5 and plotted in the middle panel of Fig. 5. Though the uncertainties are slightly larger both astrophysical parameters indicate very similar variations as in the cooler component of KR Cyg. DO Cas is also a NCB and its UBV light curves were published by Oh & Ahn (1992). We derived an effective temperature for the hotter star as 8 700 K using the intermediate-band measurements by Hilditch& Hill (1975) and the BVJHK magnitudes given in the SIMBAD database. Initial parameters were adopted from Siwak, Zola & Koziel-Wierzbowska (2010) who obtained spectroscopic mass-ratio and modelled the BV light curves. Their analysis resulted in near-contact configuration. Applying the same procedure we derive A_2 and T_2 for the irradiated star of DO Cas. The results are given in Table 5 and plotted in the bottom panels of Fig.5. The uncertainties of the parameters are very large, originated from the large scatters in the observational data, the effective albedos are smaller than the expected value of 0.5.

In Fig. 6 we plot the observed U-B and B-V colors for the systems KR Cyg, AK CMi and DO Cas. The bluer color especially in the U-B just in and around primary eclipse is clearly seen. About 50 per cent of the apparent disk of the hotter component of KR Cyg is obscured by the cooler star in mid-primary eclipse. We, therefore, suggest that multiple scattering of the light between the components of close binaries with large temperature differences results in bluer color, especially in the annular eclipse.

4. CONCLUSIONS

We obtained the five-year UBVR light curves and the radial velocities of the near-contact binary KR Cyg. Analysing the multi-color data and measured radial velocities we obtained, for the first time, the absolute parameters of the components. Comparison of the components' radii with the corresponding Roche lobes indicate that the system appears to be near-contact but not in contact. Combining the results of light-and radial-velocity analyses we obtained the absolute physical parameters of the system. Both components locate on the main-sequence of the Hertzsprung-Russell diagram. Systematic



Fig. 5. The empirical values of the effective albedos (left panels) and temperatures (right panels) are plotted against the wavelengths for the cooler components of KR Cyg (top panels), AK CMi (middle panels) and DO Cas (bottom panels).



Fig. 6. The U-B and B-V colors against the orbital phase for KR Cyg (top panels), AK CMi (middle panels) and DO Cas (bottom panels).

behaviour of the residuals between the observed and fitted light curves is attributed to the effect of the mutual heating of the components, mainly in the case of the cooler star, of this near-contact system. Analyses indicate that the effective albedo and the effective temperature of the irradiated star are significantly altered. The empirically derived value of albedo is smaller than that expected for a convective star. While the albedo gets smaller the effective temperature increases. We also analyzed the UBV data, taken from literature, for the systems AK CMi and DO Cas. The changes of albedos versus the wavelength are very similar for AK CMi and KR Cyg. **The analysis of eclipsing binary light curves involves a large number of parameters. The bolometric albedos, limb-darkening coefficients, gravity-brightening exponents are mostly kept fixed at the theoretical values. Nevertheless, external illumination is significant in close eclipsing binary systems which heats the surface layers of the illuminated star, thus alters its physical properties.**

Acknowledgment

We thank to Turkish Scientific and Technical Research Council (TÜBİTAK) for a partial support with Project number 109T708 and EBİLTEM Ege University Science Foundation Project No:2011/BİL/007.We also thank to TÜBİTAK National Observatory (TUG) for a partial support in using RTT150 and T100 telescopes with project numbers 10ARTT150-483-0, 11ARTT150-123-0 and 10CT100-101. We also thank to responsible for the Bakırlıtepe observing station for their warm hospitality. This research has been made use of the ADS and CDS databases, operated at the CDS, Strasbourg, France and TÜBİTAK ULAKBİM Süreli Yayınlar Kataloğu.

....

REFERENCES

Alencar, S. H. P. & Vaz, L. P. R., 1999, A&AS, 135, 555
Alencar, S. H. P., Vaz, L. P. R., Nordlund, A., 1999, A&A, 346, 556
Al-Naimiy, H.M.K., Al-Mahdi, H. A., Al-Sikab, A. O., Mutter, A. A., 1985, ApSS, 117, 351
Bohm-Vitense, E., 1981, An. Rev. A&A, 19, 295
Claret, A., 2001, MNRAS, 327, 989
Claret, A. & Gimenez, A., 1990, A&A, 230, 412
Cranmer, S.R., 1993, MNRAS, 263, 989
Cutri, R. M., et al., 2003, The IRSA 2MASS All-Sky Point Source Catalog, NASA/IPAC Infrared Science Archive. http://irsa.ipac.caltech.edu/applications/Gator
de Jager, C.,& Nieuwenhuijzen, H., 1987, AA, 177, 217
Drilling, J. S., & Landolt, A. U., 2000, Allen's Astrophysical Quantities, 4th ed. Edited by Arthur N. Cox. ISBN: 0-387-98746-0. Publisher: New York: AIP Press; Springer, 2000, p.381
Flower, P. J., 1996, ApJ, 469, 355

Gaposchkin, S., 1953, HA, 113, 104
Hilditch, R. W. & Hill, G., 1975, MmRAS, 79, 101
Hovhannessian, R.Kh., 2004, Astrophysics, 47, 499
Johnson, H. L. & Morgan, W. W., 1953, ApJ, 117, 313
Kirbiyik, H., 1982, MNRAS, 200, 907
Kirbiyik, H. & Smith, R. C., 1976, MNRAS, 170, 103
Lassovszky, K., 1936, AN, 258, 93
Lucy, L. B., 1967, Z. Astrophys., 65, 89
Nekrasova, C., 1945, Pulk. Bull., 134, 27
Nidever D. L., Marcy G. W., Butler R. P., Fischer D. A., and Vogt S. S., 2002, ApJS, 141, 503
Nordlund, A. & Vaz, L. P. R., 1990, A&A, 228, 231
Oh, K. D. & Ahn, Y. S., 1992, ApSS, 187, 261
Prša A., Zwitter T. 2005, ApJ, 628, 426P
Rafert, J. B. & Twigg, L. W., 1980, MNRAS, 193, 79
Rucinski, S. M., 1969, Acta Astr., 19, 245
Rucinski, S. M., 1973, AA, 23, 89
Samec, R. G., Carrigan, B. J., Gray, J. D., French, J. A., McDermith, R.J., Padgen, E. E.,1998, AJ, 116, 895
Schneller, H., 1931a, AN, 242, 180
Schneller, H., 1931b, AN, 243, 337
Schneller, H., 1931c, AN, 245, 43
Shakhovskaya, N. I., 1974, IBVS, No.897, 1
Shaw, J. S., 1990, ”in Active Close Binaries”, ed. C.Ibanoglu (Kluwer Academic Publishers, Netherlands,p.241
Shaw, J. S., 1994, MmSAI, 65, 95
Sipahi, E., Gulmen, O., 2004, ApSS, 293, 307
Sipahi, E., 2012, New Astr., 17, 383
Siwak, M., Zola, M., Koziel-Wierzbowska, D., 2010, AcA, 60, 305
Tonry, J., & Davis M., 1979, AJ 84, 1511
Topping J., 1972, ”Errors of Observation and Their Treatment”, (Chapman and Hall Ltd.), p.89
Tsesevich, V.P., 1954, Odessa Izv., 4, 263
van Hamme, W. 1993 AJ, 106, 2096
Vetesnik, M., 1965, BAC, 16, 326
Wachmann, A.A., 1948, Astr. Abh. Erg. AN, 11, No.5
Vaz, L. P. R. & Nordlund, A., 1985, A&A, 147, 281
Wilson, R. E. & Devinney, E. J., 1971, ApJ, 166, 605
Wilson, R. E. & Rafert, J. B., 1980, A&AS, 42, 195

Ege University, Science Faculty, Astronomy and Space Sciences Dept., 35100 Bornova, İzmir, Turkey. $e-mail$: esin.sipahi@mail.ege.edu.tr